

# Wildlife & Ecosystem Services: Science Objectives



**Objective #1.** To **understand** how spatial and temporal dynamics in environmental and ecological conditions within the ABoVE Study Domain influence:


(a) movement, habitat selection and population viability of a suite of highly mobile terrestrial animal species, and;

(b) accessibility of natural resources to local subsistence communities.




**Objective #2.** To **provide local stakeholders** - including natural resource agencies, wildlife managers, First Nations, Alaskan natives, and other stakeholders - **with knowledge, products, and tools** that will aid them in making informed management and adaptation decisions.





*Boulder, CO January 2017*



## **Highest priority:**

**Identify data gaps & solutions for studying  
wildlife – snow interactions**







# Integrating snow science and wildlife ecology in Arctic-boreal North America

Natalie T. Boelman<sup>1</sup>, Glen E. Liston<sup>2</sup>, Mark Hebblewhite, Laura R. Prugh<sup>Y</sup>, Arjan J. H. Meddens<sup>Z</sup>, John S. Kimball<sup>N</sup>, Peter J. Mahoney<sup>Y</sup>, Eliezer Gurarie, Anne W. Nolin, Peter B. Kirchner\*, Chris L. Cosgrove<sup>ZZ</sup>, Gil Bohrer<sup>X</sup>, Scott D. LaPoint<sup>1</sup>, Lee A. Vierling<sup>Z</sup>, Jan U.H. Eitel<sup>Z</sup>, Todd J. Brinkman<sup>3</sup>, Stine Højlund Pedersen<sup>2,XX</sup>

**Goal:** To demonstrate the need for temporally and spatially dynamic snow products to help understand how changing snowscapes impact wildlife in ABRs.

<sup>1</sup> Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY 10964, USA

<sup>2</sup> Cooperative Institute for Research in the Atmosphere, Colorado State University, Fort Collins, CO 80523, USA

<sup>sdl</sup> Department of Migration and Immuno-ecology, Max-Planck Institute for Ornithology, Radolfzell 78315, Germany

<sup>X</sup> Department of Civil, Environmental and Geodetic Engineering, The Ohio State University, Columbus, OH 43210, USA

<sup>N</sup> Numerical Terradynamic Simulation Group, W. A. Franke College of Forestry and Conservation, University of Montana, Missoula, MT 59812, USA

<sup>XX</sup> Department of Biological Sciences, University of Alaska Anchorage, Anchorage, AK 99508, USA

<sup>Y</sup> School of Environmental and Forest Sciences, University of Washington, Seattle WA 98195

<sup>Z</sup> Department of Natural Resources and Society, University of Idaho, Moscow, ID 83844, USA

\* Southwest Alaska Network, National Park Service, Anchorage, AK 99501, USA

<sup>3</sup> Institute<sup>ZZ</sup> College of Arctic Biology, Earth, Ocean and Atmospheric Sciences, Oregon State University of Alaska Fairbanks, Fairbanks, AK 99775, USA, OR 97331

in preparation for *Environmental Research Letters*, ABoVE special issue

# I. INTRODUCTION

- snow covered 9-10 months/year
  - “Snow has it’s fingers in everything” – *Matthew Sturm*
- snowscapes have been changing in recent decades
- ✓ consequences on biogeochemistry, hydrology & energy balance
- ? consequences on wildlife
- one of the last remaining regions of the planet with intact wildlife communities
- yet the snow data available are often unsuitable/not the most critical:
  - spatial & temporal resolutions
  - spatial & temporal extents
  - physical snowpack variables

**If I’d known the  
SWE was so high  
over here, I would  
have stayed away!**





## II. THE IMPORTANCE OF SNOWSCAPES TO WILDLIFE



Timing of egg laying

Liebezeit et al. 2014; Green et al. [1977](#); Meltofte et al. [2007](#); Smith et al. [2010](#); Grabowski et al. [2013](#); Boelman et al. 2017



Predator-prey dynamics




Berteaux et al. 2016; Bilodeau et al. 2013; Duchesne et al. 2011

Zimova et al. 2014; Mills et al. 2013; Henden et al. 2017



## II. THE IMPORTANCE OF SNOWSCAPES TO WILDLIFE



Trophic cascade



Fancy and White 1987; Nicholson et al. 2016; Verme 1968; Kelsall and Telfer 1971; Mech et al. 1971; Kelsall and Prescott 1971; Telfer and Kelsall 1984; Pimlott et al. 1969; Mech and Frenzel 1971; Haber 1977; Mech and Karns 1977; Peterson 1977; Eide and Ballard 1982; Peterson and Allen 1974; Haber 1977; Gasawy et al. 1983; Lendrum et al. 2017, Formozov 1946; Nelson and Mech 1986; Hoefs and McTaggart-Cowan 1980; Duquette 1988; Nichels and Bunnell 1999; Johnson et al. 2001; Beumer 2017, Johnson et al. 2001; Putkonen et al. 2009; Rennert et al. 2009; Stien et al. 2012; Hansen et al. 2011; Hansen 2013; Sokolov et al. 2016



## II. THE IMPORTANCE OF SNOWSCAPES TO WILDLIFE



- Depth
- Density
- Ice layers



**Insulative  
capacity**


Craighead and Craighead 1972; Jonkel 1980; Vroom et al. 1980

**A broad suite of snowscape characteristics must be considered!**

# III. CURRENT GEOSPATIAL SNOWSCAPE PRODUCTS FOR ABRs



*in-situ* measurements



remotely sensed observations



numerical modeling

model-data assimilation & reanalysis products

All summarized in a comprehensive table – thank you Arjan Meddens!



# III. CURRENT GEOSPATIAL SNOWSCAPE PRODUCTS FOR ABRs



*in-situ* measurements

**#2**


Many sites in the ABR measure:

- SWE
- snow depth
- air temperature

but rarely measure other variables

- ice layers
- hardness

# III. CURRENT GEOSPATIAL SNOWSCAPE PRODUCTS FOR ABRs



**Daily, vast coverage & spatially continuous** snow products:

- albedo
- grain size
- relative ice & water content

**But...**

**#1** most wildlife relevant snowscape properties not available

**#2** tradeoff between spatial & temporal resolutions

**remotely sensed observations**



# III. CURRENT GEOSPATIAL SNOWSCAPE PRODUCTS FOR ABRs



numerical modeling

**Can be tailored to fit the specific needs of wildlife studies!**

- produce a large suite of wildlife-relevant snow variables
- over large spatial and temporal extents
- over a range of spatial and temporal resolutions

**But, the majority are not yet up to the task:**

**#1** most lack adequate spatial resolution and the ability to run at different spatial resolutions (e.g. meters to kilometers)

**#2** most do not simulate wildlife relevant snowpack properties because:

- only as good as meteorological forcing inputs
- difficult to develop snow models that are general enough to do a good job everywhere

# IV. THE IMPORTANCE OF WILDLIFE-RELEVANT SNOWSCAPE CHARACTERISTICS AT APPROPRIATE ANALYSIS SCALES



**Case Study 1**




**Case Study 2**




**Case Study 3**



# Case study 1:




- physically based numerical model (SnowDens-3D) to map polar bear snowdrift den habitat
- simulated the year-specific physical interactions of snow, wind, & topographic dynamics (2.5 m<sup>2</sup> grid)
- created annual maps of snow depth → potential den locations (1995 – 2012)




**97% of observed den locations  
correctly identified by SnowDens-3D**

*from Liston et al. 2016*

# Case study 1: Mapping polar bear den habitat requires accurate & fine-scale snow depth estimates



- physically based numerical model (SnowDens-3D) to map polar bear snowdrift den habitat
- simulated the year-specific physical interactions of snow, wind, & topographic dynamics (2.5 m<sup>2</sup> grid)
- created annual maps of snow depth → potential den locations (1995 – 2012)



**97% of observed den locations  
correctly identified by SnowDens-3D**

*from Liston et al. 2016*

## Case study 2:



**Laura Prugh:** Assessing Alpine Ecosystem Vulnerability to Environmental Change Using Dall Sheep as an Iconic Indicator Species

Evaluated the efficacy of:


1. MODIS snow-cover fraction
2. SnowModel (Liston and Elder 2006) snow depth & density products

to predict Dall sheep movements at multiple spatial & temporal scales.

*from Mahoney et al. 2017, in revision*





## Case study 2:



# Case study 2: : Understanding Dall sheep movement behavior requires fine- and coarse-resolution snowscape products, including snow depth

Best  
model fit

Worst  
model fit



- at fine scales, Dall sheep generally selected for:

## (1) low density, shallow snow

- likely to facilitate access to forage and reduce energy expenditure

## (2) higher snow density, deep snow


- reduce snow hoof penetration & improve efficiency of movement

from Mahoney et al. 2017, in revision

# Case Study 3: To determine if the timing of caribou spring migration in NW Canada is related to inter-annual variation in snowmelt date as estimated from MODIS snow cover fraction.

**No relationship between snowmelt timing & onset of migration over 14 year period**


(supports what Le Corre et al. 2017 in NE Canada)



Elie Gurarie *in prep.*



# Case Study 3:



## Case Study 3:

- timing of spring migration *not* linked to timing of snow cover melt
- but animals prefer to migrate over complete snow cover
- Le Corre et al. (2017) suggest they are 'chasing' high *quality* snow cover






# V. A FUTURE PROSPECTUS FOR IMPROVING WILDLIFE-SPECIFIC SNOWSCAPE PRODUCTS FOR ABRs



*In situ* measurements

*Limitations  
&  
Missing data*



remotely sensed observations

*Limitations  
&  
Missing data*



numerical modeling


# V. A FUTURE PROSPECTUS FOR IMPROVING WILDLIFE-SPECIFIC SNOWSCAPE PRODUCTS FOR ABRs



## ***In situ* measurements**

Detailed temporal coverage

*Limitations  
&  
Missing data*



## **remotely sensed observations**

Vast spatial coverage

*Limitations  
&  
Missing data*



## **numerical modeling**

Large suite of wildlife-relevant variables

## **Data-model fusion**

A synergetic extraction and dynamic merging of information obtained from different sources that can 'fill in' for one another.


Glen Liston



# Snow Depth: Arctic National Wildlife Refuge (ANWR)

*in situ* Observations

(spatially interpolated)



# VI. CONCLUSIONS

- We need fit-for-purpose snow products in ABRs
- We hope that ABoVE will lead the way in improving snow science
- We think Data-model fusion is the the way to go
- But this requires active development & contributions from all 3 constituents







# OUR NEXT SYNTHESIS ACTIVITIES ?

- changing seasonality & veg. phenology

vs.

wildlife phenology & seasonality in subsistence resource availability

- wildlife – fire interactions



# Wildlife & Ecosystem Services Working Group

Boelman, Natalie – Lamont-Doherty Earth Observatory, Columbia University

Bohrer, Gil -- Ohio State University

Brinkman, Todd -- University of Alaska, Fairbanks

Chen, Wenjun -- Canada Centre for Mapping and Earth Observation

Clark, Karin -- GNWT, ENR, Wildlife

Cold, Helen -- University of Alaska, Fairbanks

Cosgrove, Christopher -- Oregon State University

Fienup-Riordan, Ann -- Calista Education and Culture, Inc.

Frost, Gerald (JJ) -- Alaska Biological Research, Inc

Gill, Michael (Mike) -- Polar Knowledge Canada

Goetz, Scott -- Northern Arizona University

Griffith, Peter -- NASA GSFC / SSAI

Gurarie, Eliezer -- University of Maryland

Hebblewhite, Mark -- University of Montana

Kimball, John -- University of Montana

Kirchner, Peter -- National Park Service

Scott LaPoint -- Lamont-Doherty Earth Observatory, Columbia Univ.

Macander, Matthew -- Alaska Biological Research, Inc.

McCaffery, Brian -- Fish and Wildlife Service

Meddens, Arjan -- University of Idaho

Miller, Charles (Chip) -- NASA JPL

Nolin, Anne -- Oregon State University

Oliver, Ruth -- Columbia University

Prugh, Laura -- University Of Washington

Reynolds, Joel -- U.S. Fish and Wildlife Service

Sowl, Kristine -- USFWS Yukon Delta National Wildlife Refuge

Vierling, Lee -- University of Idaho



H.T. HARVEY & ASSOCIATES  
Ecological Consultants

